# Miami-Dade Urban Tree Canopy Analysis

Henry Hochmair Associate Professor Geomatics Program University of Florida Daniel Gann
Research Associate
GIS-RS Center
Florida International University

Adam Benjamin Geomatics Specialist Geomatics Program University of Florida Jennifer Fu Head of GIS-RS Center Florida International University







# **Funding Agencies**

 Miami-Dade County Parks and Recreation - Neat Streets Miami







American Forests



#### **Project Scope**


- Determine for the Urban Area of Miami-Dade County:
  - areas with current tree canopy (existing UTC)
  - areas with potential tree canopy (possible UTC)
  - areas unsuitable for tree canopy





#### **Project Scope**


- Determine for the Urban Area of Miami-Dade County:
  - areas with current tree canopy (existing UTC)
  - with potential tree canopy (possible UTC)
  - unsuitable for tree canopy
- Determine the UTC percentage breakdown for
  - census places
  - commission districts
  - land use categories
  - property parcels





# **Project Scope**

- Determine for the Urban Area of Miami-Dade County:
  - areas with current tree canopy (existing UTC)
  - with potential tree canopy (possible UTC)
  - unsuitable for tree canopy
- Provide the percentage breakdown for
  - census places
  - commissioner districts
  - property parcels
  - selected land use categories
- Relate percent of current UTC to
  - surface temperature
  - socio-economic data





#### **Study Area**

- Urban Area of Miami-Dade County
- 1150 km² (444 mi²)






#### **Method**

#### Create a land cover classification map, using:

- Satellite imagery
  - WorldView II: 8 band spectral resolution, 2m spatial resolution





#### **Method**

#### Create a land cover classification map, using:

- Satellite imagery
  - WorldView II: 8 band spectral resolution, 2m spatial resolution
- GIS vector layers from County and other sources, e.g.
  - Large and small buildings
  - Edge of pavement, railroads
  - Water bodies
  - Agricultural areas





#### **Method**

#### Create a land cover classification map, using:


- Satellite imagery
  - WorldView II: 8 band spectral resolution, 2m spatial resolution
- GIS vector layers from County and other sources, e.g.
  - Large and small buildings
  - Edge of pavement, railroads
  - Water bodies
  - Agricultural areas
- Apply GIS spatial analysis functions for areal statistics






# **Results: Land Cover Classification Map**

9 land cover categories





#### **Land Cover Statistics**




| Class           | Area (km²)       | Percent cover  |  |
|-----------------|------------------|----------------|--|
| Tree Canopy     | 228.6 ( ± 13.7 ) | 19.9 ( ± 1.2 ) |  |
| Street/Railroad | 117.2 ( ± 4.4 )  | 10.2 ( ± 0.4 ) |  |
| Building        | 183.1 ( ± 11.7 ) | 15.9 ( ± 1 )   |  |
| Wetland         | 18.7 ( ± 4.2 )   | 1.6 ( ± 0.4 )  |  |
| Water           | 58.6 ( ± 4.6 )   | 5.1 ( ± 0.4 )  |  |
| Cropland        | 35.6 ( ± 0.9 )   | 3.1 ( ± 0.1 )  |  |
| Grass           | 255.8 ( ± 15.6 ) | 22.2 ( ± 1.4 ) |  |
| Bare Ground     | 21.4 ( ± 0.9 )   | 1.9 ( ± 0.1 )  |  |
| Impervious      | 230.9 ( ± 13.5 ) | 20.1 ( ± 1.2 ) |  |
| Total           | 1150.0           | 100.00%        |  |



#### **Land Cover Statistics**


- Total possible tree canopy: 44.2%
- Possible tree canopy on pervious surfaces: 24.1%





# **Tree Canopy and Surface Temperature**

Temperature in Degree Fahrenheit



Landsat Enhanced Thematic Mapper (ETM) thermal band (30m)





# **Tree Canopy and Surface Temperature**

Tree Canopy

Low: 58.6478

1,000

1,000


2,000

2,000

Feet

Heat map with tree canopy






Contour map



# Land Cover Mix and Temperature (°F)



Bivariate correlations between percent land cover and surface temperature:

|   | % Impervious | % Street/RR | % Tree | % Water |
|---|--------------|-------------|--------|---------|
| r | 0.410        | 0.346       | -0.432 | -0.171  |
| р | < 0.001      | <0.001      | <0.001 | < 0.001 |




# **Analysis of Census Places**




FLORIDA INTERNATIONAL UNIVERSIT

79 census places




# **Analysis of Census Places**






# **Analysis of Census Places: Subset**






#### **Analysis of Census Places: Maps**

#### % Existing UTC



#### % Possible UTC (Pervious)





17.0%

#### **Indian Creek**

• 68.6 % Possible UTC (Pervious)









# **Existing and Possible Canopy in Parcels**
















# UF FLORIDA


#### **Analysis of Socioeconomic Variables**

- Analysis for 1525 populated census block groups
- Relate % tree canopy to
  - median annual household income in US \$
  - percent African American population
  - percent Hispanic population




# UF | UNIVERSITY of FLORIDA

# **Analysis of Socioeconomic Variables: Maps**



# **Bivariate Correlations with % Tree Canopy**



# **Summary**

- Overall tree canopy was 19.9%
- Residential housing represents 42% of the existing tree canopy and 33% of possible tree canopy on pervious surfaces
- Tree canopy and water bodies are associated with lower surface temperatures
- Tree canopy is positively correlated with median income, but negatively correlated with percentage of African American and Hispanic residents

